



# Gooseberries & Currants

Cheryl Kaiser<sup>1</sup> and Matt Ernst<sup>2</sup>

## Introduction

Gooseberries and currants (*Ribes* spp.) are woody, multi-stemmed shrubs best known for their tart fruit. While some enjoy eating them fresh, these fruit are especially prized for use in making jellies, jams, pies, and sauces.

## Marketing

Gooseberries and currants are marketed fresh, preserved (frozen and canned) and in value-added food items, like fruit preserves and baked goods. Market-place options include farmers markets, on-farm stands and other direct markets. Upscale or “white tablecloth” restaurants that make their own sauces and desserts may also be a market possibility for gooseberries and currants. Gooseberries and currants can diversify the product mix at a farmers market, roadside stand, and CSA (community supported agriculture). Providing consumer education through suggested uses, recipes, and other “point-of-purchase” materials may help producers develop a more steady market for gooseberries and currants.

## Market Outlook

Currants and gooseberries are minor crops in the U.S., with fewer than 2,500 commercial acres, according to the U.S. Census of Agriculture. Historically, the U.S. regulated cultivation of *Ribes* species in an effort to control white pine blister rust (WPBR) mortality in pines; the WPBR fungus needs a *Ribes* plant to complete its life cycle. The federal restriction was lifted in 1966, but some states continue to restrict gooseberry and currant production. Production of *Ribes* is not restricted in Kentucky, where white pine blister rust appears to be uncommon.



GOOSEBERRIES

Producers should estimate both local production and demand before investing in producing and marketing gooseberries and currants. Preserving the berries for sale during the off-season may also be a way of diversifying a grower’s offerings and spreading the risk of production for a new crop across different target markets. Evaluations of *Ribes* cultivars for Kentucky have been made by Kentucky State University researchers, who reported market potential in Kentucky for white, red, and black currants, as well as gooseberries. Gooseberries have the greatest potential for fresh market sales. However, gooseberries also are relatively easy to grow and local supply can quickly meet consumer demand.

## Production Considerations

*Crop/cultivar selection and propagation*

Currant types are red, white, and black; American and European gooseberries are green to yellow or red when ripe. Cultivars vary in terms of fruit characteristics and disease resistance. Consort,



<sup>1</sup>Cheryl Kaiser is a former Extension Associate with the Center for Crop Diversification.

<sup>2</sup>Matt Ernst is an independent contractor with the Center for Crop Diversification.

Coronet, Crusader, and Titania are immune to WPBR. Based on Kentucky State University research trials, the cultivars listed in TABLE 1 performed well in Kentucky. Selections with disease susceptibility are also included because they were superior in other production traits to those with resistance. WPBR susceptible cultivars should not be planted within 1,000 to 3,000 feet of susceptible pines.

Currants are easily propagated by hardwood cuttings of 1-year-old wood, while tip layering is a surer method of gooseberry propagation.

*Site selection and planting*

Gooseberries and currants prefer a rich, moist, well-drained soil; however, they will tolerate a wide range of soil types. Avoid waterlogged soils, as well as low sites where spring frosts might settle and injure blossoms. Northern exposures that offer some afternoon shade are preferred, as is good air circulation.

Site preparation, including the destruction of all weeds, a year prior to planting is recommended. Well-rooted 1-year-old cuttings are planted during dormancy in late fall or early spring. Black currant cultivars are generally self-sterile, thus cross pollination is necessary. Gooseberries, red currants, and white currants, on the other hand, are self-fruitful so multiple cultivars are not required for pollination; however higher yields and larger fruits can be obtained with cross-pollination.



WHITE CURRANTS (ABOVE) AND RED CURRANTS (BELOW)



The majority of fruit is produced on the 2- and 3-year-old shoots; canes that are 4 years or older produce very little fruit. Mature plants should be pruned annually during dormancy to remove the older canes and

Table 1. GOOSEBERRY AND CURRANT CULTIVARS THAT HAVE PERFORMED WELL IN KENTUCKY

|                      | Resistant to WPBR <sup>1</sup> and PM <sup>2</sup> | Resistant to WPBR; not PM | Resistant to PM; not WPBR | Not resistant to WPBR nor PM |
|----------------------|--|---------------------------|---------------------------|------------------------------|
| <b>Gooseberries</b>  | Hinnomaki Red<br>Amish Red<br>Poorman              |                           |                           |                              |
| <b>Black currant</b> | Titania  |                           | Ben Sarek<br>Ben Lomond   |                              |
| <b>Red currant</b>   | Viking   | Jonkeer Van Tets          |                           | Rovada                       |
| <b>White currant</b> | Primus   |                           | White Imperial            |                              |

<sup>1</sup>WPBR = White Pine Blister Rust    <sup>2</sup>PM=Powdery Mildew

to promote new growth. Pruning helps increase yields and keeps the bushes manageable. Irrigation in June and July is important for improved fruit size and reducing heat stress.

#### *Pest management*

It is recommended that growers generally select varieties that have white pine blister rust and powdery mildew resistance. Other potential disease problems include Septoria leaf spot and anthracnose.

While these crops have few serious insect pests, injury from mites and aphids can occur occasionally. Other potential pests include currant fruit fly and currant borer. The imported currantworm is a sawfly larva that can rapidly strip the foliage from plants. Watch for this greenish-gray larva covered with dark spots beginning just after the leaves emerge in the spring. In some seasons birds can be a serious problem as fruit ripens. Good weed control is very important and can be accomplished with cultivation and/or the use of mulch. Kentucky growers may be able to use organic methods in producing these crops.

#### *Harvest and storage*

Gooseberries and currants ripen during mid- to late June in Central Kentucky and have roughly a two-week harvest period. Fruit for processing is generally hand-picked when fully sized but not completely ripe. Unripe currants can be placed in cold storage for a few days, if necessary, while gooseberries can keep for up to two weeks. Fruit for fresh use should be picked when fully ripe and marketed promptly.

There is a time lapse of three years after planting before the first harvest is realized, with full production at four years. Unlike currants, gooseberries have needle-like thorns that slow harvest. Mature bushes can yield 4 to 8 pounds of fruit. Properly managed gooseberries and currants can continue to fruit for 15 years or more.

#### *Labor requirements*

Because of limited market demand, it is unlikely that gooseberries and currants will be planted in Kentucky on a per-acre basis. Labor needs per 120 plants (about one-fifth acre) are approximately 10 hours for production and 40 hours for harvest. Additional labor time will be required for processing these crops.

### **Economic Considerations**

Initial investments include land preparation, plant es-



GOOSEBERRIES

tablishment, and installation of an irrigation system. The most significant investment for gooseberries and currants will be the initial plant purchase, as plants can be difficult to obtain and should be ordered well in advance. Producers should be prepared to pay up to the \$7 range per plant. Assuming a price of \$3.50 per pound (about \$5 to \$5.50 per quart), establishment costs will not be fully paid back until Year 5 of production.

Establishment costs for a one-fifth acre planting are estimated in 2019 at \$3,860 for the first three years, including about \$1,300 in labor costs. In a full-bearing year, production costs are estimated at \$300 per one-fifth acre, with harvest and marketing costs at about \$900. Total costs per one-fifth acre in a full-production year, including both variable and fixed, were estimated between \$1,800 and \$1,900 for 2019. Presuming gross returns (conservative) of \$2,500 from full production per one-fifth acre, returns above variable costs could be approximately \$1,000 per one-fifth acre with a return to land, capital and management of \$750 to \$800 per one-fifth acre. Like other properly marketed small fruit, gooseberries and currants have the potential for net returns in the \$6,000 to \$8,000 range per acre, especially through direct markets; however, Kentucky's current local demand is not likely to support widespread commercial plantings of *Ribes*.

### **Selected Resources**

#### *On the Internet*

- Growing Gooseberries and Currants (Kentucky State University, 2010) 3.84 MB Powerpoint <http://kysu.edu/wp-content/uploads/2017/08/goosetalk02112010fin2000.pdf>

- Currant and Gooseberry Genetic Resources (USDA-ARS National Clonal Germplasm Repository, 2010) <http://www.ars.usda.gov/Main/docs.htm?docid=11353>
- Currants and Gooseberries (Purdue University, 2001) [https://edustore.purdue.edu/item.asp?item\\_number=ho-17-w#.WECJUpIwbxU](https://edustore.purdue.edu/item.asp?item_number=ho-17-w#.WECJUpIwbxU)
- Currants and Gooseberries (Ohio State University, 2011) <https://southcenters.osu.edu/horticulture/fruits/currants-and-gooseberries>
- Gooseberry (Agricultural Marketing Resource Center, 2016) <https://www.agmrc.org/commodities-products/fruits/gooseberry>
- Gooseberries and Currants (Cornell University, 2019) <http://www.hort.cornell.edu/fruit/mfruit/gooseberries.html>
- The Berry Diagnostic Tool (Cornell University) <https://blogs.cornell.edu/berrytool/>

- Specialty Crop Profile: Ribes (Gooseberries and Currants) (Virginia Cooperative Extension, 2009) <https://www.pubs.ext.vt.edu/438/438-107/438-107.html>

*In print*

- Currants, Gooseberries, and Jostaberries: A Guide for Growers, Marketers, and Researchers in North America. Danny L. Barney and Kim E. Hummer. 2005. Hawthorn Press. 266 pp.

**Suggested Citation:**

Kaiser, C. and M. Ernst. (2020). *Gooseberries & Currants*. CCD-CP-6. Lexington, KY: Center for Crop Diversification, University of Kentucky College of Agriculture, Food and Environment. Available: <http://www.uky.edu/ccd/sites/www.uky.edu/ccd/files/currants.pdf>

---

*Reviewed by John Strang, UK Extension Specialist, and Sheri Crabtree, Kentucky State University Horticulture Research and Extension Associate*

*Photos courtesy of [Pixabay.com](http://Pixabay.com)*

**September 2020**

---

For additional information, contact your local **County Extension agent**

Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, or physical or mental disability.